

# *Journal of Theoretics*

Volume 6-3, June/July 2004

---

## **A New Equation of Light Trajectory**

Dipl. Ing. Andrija S. Radović

Address: Nike Strugara 13a, 11030 Beograd, Yugoslavia

[andrijaradovic@hotmail.com](mailto:andrijaradovic@hotmail.com)

**Abstract:** The article shows that the essential equation of light trajectory is very simple and that is connected with the third derivative of the coordinate oppositely than the classic mechanics where the second derivative is the last significant one.

**Keywords:** light, trajectory, deflection.

### ***Introduction***

Light movement can be well described only by its wave nature and formula for the refraction of light between two media. The article shows that every localized carrier of the light has trajectory described with the equation (9). The solution of equation (9) is especially interesting when  $v$  is constant.


### ***Derivation***

Let start from the well-known equation of light refraction:

$$\frac{\sin(\theta_1)}{\sin(\theta_2)} = \frac{v_1}{v_2} \quad (1)$$

Whereas:

- $v_1$  = the speed of light in medium 1,
- $v_2$  = the speed of light in medium 2,
- $\theta_1$  = angle of light's deflection in medium 1,
- $\theta_2$  = angle of light's deflection in medium 2.




We can suppose that the mediums are nearly equals and that the light speeds are also nearly equals, than we have:

$$\frac{\text{SIN}(\theta + d\theta)}{\text{SIN}(\theta)} = \frac{v + dv}{v} \quad (2)$$

⇒

$$\frac{\text{SIN}(\theta) \cdot \text{COS}(d\theta) + \text{COS}(\theta) \cdot \text{SIN}(d\theta)}{\text{SIN}(\theta)} = 1 + \frac{dv}{v} \quad (3)$$



It can be derived directly:

$$\frac{\text{COS}(\theta)}{\text{SIN}(\theta)} \cdot d\theta = \frac{dv}{v} \quad (4)$$

⇒

$$\frac{\text{COS}(\theta)}{\text{SIN}(\theta)} = \frac{1}{v} \cdot \frac{dv}{d\theta} \quad (5)$$

⇒

$$\text{TAN}(\theta) = \frac{v}{\left(\frac{dv}{d\theta}\right)} \quad (6)$$

⇒

$$\theta = \text{ATAN}\left(\frac{v}{\left(\frac{dv}{d\theta}\right)}\right) \quad (7)$$

After the both sides of equation are differentiated by  $\theta$ , hence the following formula is obtained:

$$v \cdot \frac{d^2v}{d\theta^2} + v^2 = 0 \quad (8)$$

Finally:

$$\boxed{v + \frac{d^2 v}{d\theta^2} = 0} \quad (9)$$

I.e.

$$|\vec{v}| + \frac{d^2 |\vec{v}|}{d\theta^2} = 0 \quad (10)$$

This is the final equation of the light trajectory, which gives the basic relation between the path of the light beam and its differential declination. Evaluated equation for  $E^3$  space is given by the following equation:

$$\vec{v}^2 \cdot (\vec{a} \times \dot{\vec{a}}) = 2 \cdot \vec{a}^2 \cdot (\vec{v} \times \vec{a}) \quad (11)$$

## Conclusion

This is the final light beam trajectory equation. It shows that there is a great influence of the third time derivation of the coordinates, while in classic mechanics the second derivative is the last significant one. It also shows that light changes its speed near the rigid bodies' edges and in gaps too. The equation could be used for testing purposes in new electromagnetic theories.

## References

1. Siepmann JP, "The Light Clock: A New Method of Measuring True Time", Journal of Theoretics.
2. Ioannis Iraklis Haranas, "A Simple Calculation of a Possible Variation in the Speed of Light", Journal of Theoretics.
3. V.N. Strel'tsov, "Light Velocity in General Relativity", Journal of Theoretics Vol.4-5.
4. William Rowan Hamilton, "ON A GENERAL METHOD OF EXPRESSING THE PATHS OF LIGHT, AND OF THE PLANETS, BY THE COEFFICIENTS OF A CHARACTERISTIC FUNCTION", Dublin University Review and Quarterly Magazine, 1 (1833), pp. 795–826.
5. William Rowan Hamilton, "ON A MODE OF DEDUCING THE EQUATION OF FRESNEL'S WAVE", Philosophical Magazine, 19 (1841), pp. 381–383.
6. William Rowan Hamilton, "ON SOME RESULTS OF THE VIEW OF A CHARACTERISTIC FUNCTION IN OPTICS", British Association Report, Cambridge 1833, pp. 360–370.

Received May 2003

[Journal Home Page](#)

© Journal of Theoretics, Inc. 2004